

Diagrams of Troosts Crinoides.


Actinocrinus gibboous Troost



actinocrinus comutus Front





Actinocrinees wind Troost.




actinocrinus Tennessee, Troot. = S'accocrimus speciosus, hoemer. not of Hall. actinocrims fibula, Troot. actinoerius Manneri, Front, Very like a. cornetus, J. but with a more spreading Calyx. Cichnocrinus - sp? Hall in Drovst This specimen was probaced under act. mortoni by Troost but it differs from his description in the sumber of limes having only four to each very while that species has six

actinocrims Lea. Twost = Conocrincles Lea, Frost. = Aloprosalo crimis comens, Lyon; Converintes Leai? Drost of ______ as given by him to Mr White among some fossils sent to Mr Hall from Crawfording by Rol. Hovey Cacabocrinus sculptus Turst.








.




Pentremites Neimwardti Troost. No1468





























actinocrims Lea. Troost = Convermétés Lea, Groot. = Aloprosalo crimis comens, Lyon; as given toy him to Mr White among some fossils sent to Mr Hall from Crawfording, by Prof. Hovey Cacabocrinus sculptus Turst.









Encaly ptocrims splendidus Twost.



Encalyptocrims goldfussi Troost-= E. Phillippsii var. Hall in Droost




Balanocrims sculptus, Twost,

-Lampterocrims Dennesseae, Twomer,

The five divisions of the basal


plates were taken from

Apecimens in Prof Halls, Coll.





Cyathocrims stellatus, Twost. Ey athornins demessee, Troost, Egathorius pentaspheneus, Frost, Egathocrims corrugatus, Front very like 6. sculptiles, Heell: from Burling ton limes tom











- Donacicrimites simplex, broost; in M.S. Catillocrinus Dennessaue, Twost.

